

Gauge Theories Coupled to Fermions in Generation

Hiromi KASE, Katsusada MORITA* and Yoshitaka OKUMURA**

Department of Physics, Daido Institute of Technology, Nagoya, 457

**Department of Physics, Nagoya University, Nagoya, 464-01*

***Department of Natural Sciences, Chubu University, Kasugai, Aichi, 487*

Gauge theories coupled to fermions in generation are reformulated in a modified version of extended differential geometry with the symbol χ . After discussing several toy models, we will reformulate in our framework the standard model based on Connes' real structure. It is shown that for the most general bosonic lagrangin which is required to also reconstruct $N = 2$ super Yang-Mills theory Higgs mechanism operates only for more than one generation as first pointed out by Connes and Lott.

§1. Introduction

An ingenious way of 'geometrizing' Higgs mechanism, namely, unifying gauge and Higgs fields via Dirac operator on a product space of Minkowski space-time with two-points internal space was proposed by Connes^{1), 2), 3), 4)} who applied his noncommutative geometry (NCG) to reconstruct the standard model. The gauge group is defined to be the unitary group of a noncommutative algebra which is represented on Hilbert space of fermions. The algebra representation (rep) naturally leads to the concept of generation. It is also remarkable that pure Yang-Mills action functional in NCG automatically contains Higgs potential.

NCG approach initiated by Connes has been followed by many works^{5), 6), 7), 8), 9), 10), 11)} in the same or similar veins. Among them Sitarz⁹⁾ first pointed out that NCG admits a differential geometric formulation in terms of the symbol χ^* , which was forced to satisfy curious properties. For instance, unlike dx^μ , χ is not closed, $d_\chi \chi \neq 0$ and the exterior product is symmetric, $\chi \wedge \chi \neq 0$. In fact, it is possible to define^{12), 13)} a consistent noncommutative differential geometry with $d_\chi \chi = 0$ for model building provided $\chi \wedge \chi \neq 0$.

On the other hand, it was shown by Chamseddine¹⁴⁾ that NCG is capable to reconstruct $N = 2, 4$ super Yang-Mills theories. We have notified¹⁵⁾, however, that to describe such theories using χ we have to abandon the symmetry $\chi \wedge \chi \neq 0$ and adopt the new relation $\chi \wedge \chi = 0$ with the field strength redefined. This lets χ become more similar to the usual one-form basis¹⁶⁾.

Connes^{4), 5)} also introduced the real structure into NCG. This concept has been discussed more explanatorily in several related works^{17), 18), 19)}. It means that flavor and color symmetries are to be implemented in the bimodule structure of the Hilbert space.

The purpose of this paper is first to present a modified version of extended differential geometry¹⁶⁾, next to apply it to broken gauge theories coupled to fermions in generation and third to discuss the standard model in our framework relating the real structure with the factorization property of the standard model gauge transformations.

The next section defines a modified form of an extended differential geometry. Toy models will be discussed in Section 3. The standard model will be treated in the section 4. The section

^{*)} The symbol χ constitutes the fifth one-form basis in addition to the usual four one-form basis.

5 is devoted to discussions. Some calculational details are postponed to the Appendix.

§2. Extended differential geometry

Following Connes we shall consider an unital, involutive noncommutative algebra $\mathcal{A}$ together with its $*$ -preserving rep ρ acting on the Hilbert space $\mathcal{H}$ of fermions such that

$$\rho(ab) = \rho(a)\rho(b), \quad \rho(a+b) = \rho(a) + \rho(b), \quad \rho(1) = 1, \quad \rho(a^*) = \rho(a)^\dagger \quad (2.1)$$

where $a, b \in \mathcal{A}$, $a \mapsto a^*$ is the involution and † denotes hermitian conjugation. In NCG $\rho(\mathcal{A})$ is the set of multiplicative operators on the fermion Hilbert space $\mathcal{H}$, $(a, \psi) \mapsto \rho(a)\psi$, $a \in \mathcal{A}, \psi \in \mathcal{H}$.

For simplicity we omit the notation ρ in the rest of this section. The massless Dirac lagrangian takes the form

$$\mathcal{L}_D = i\bar{\psi}\gamma \cdot \partial\psi \equiv i\langle\tilde{\psi}, d\psi\rangle, \quad \tilde{\psi} = \gamma_\mu\psi d\hat{x}^\mu, \quad \langle d\hat{x}^\mu, d\hat{x}^\nu \rangle = g^{\mu\nu}, \quad g^{\mu\nu} = \text{diag}(1, -1, -1, -1) \quad (2.2)$$

d denoting the usual exterior derivative in ordinary differential geometry, $d\psi = \partial_\mu\psi d\hat{x}^\mu$, and hat meaning dimensionless. Clearly this is not invariant under the gauge transformation

$$\psi \xrightarrow{g} \psi = g\psi, \quad g \in \mathcal{U}(\mathcal{A}) \quad (2.3)$$

where the unitary group of the algebra $\mathcal{A}^*$)

$$\mathcal{U}(\mathcal{A}) = \{g \in \mathcal{A}; gg^* = g^*g = 1\} \quad (2.4)$$

defines the gauge group. The lack of gauge invariance is remedied by replacing $i\langle\tilde{\psi}, d\psi\rangle$ with

$$\mathcal{L}_D = i\sum_i \langle a^{i*}\tilde{\psi}, d(b^i\psi) \rangle, \quad a^i, b^i \in \mathcal{A} \quad (2.5)$$

and assuming in addition to Eq.(2.3) the following gauge transformation^{**)}

$$a^i \rightarrow ga^i, b^i \rightarrow b^ig^*, g \in \mathcal{U}(\mathcal{A}) \quad (2.6)$$

We assume that $\sum_i \langle a^{i*}\tilde{\psi}, b^i\psi \rangle = \langle \tilde{\psi}, \psi \rangle$, which implies the relation $\sum_i a^ib^i = 1$ ^{7)***)}.

Equation (2.5) is re-written as

$$\mathcal{L}_D = i\langle\tilde{\psi}, (d + A)\psi\rangle, \quad (2.7)$$

where we have introduced antihermitian, differential one-form

$$A = \sum_i a^i db^i = A_\mu d\hat{x}^\mu, \quad a^i, b^i \in \mathcal{A} \quad (2.8)$$

^{*)} The algebra $\mathcal{A}$ contains as a subset the commutative algebra $C^\infty(M_4)$ defined over Minkowski space-time M_4 .

^{**)} We make use of the equivalence relation $(ag^*, g\psi) \sim (a, \psi)$, $g \in \mathcal{U}(\mathcal{A})$.

^{***)} This requires a (finite) sum in Eq.(2.5).

Equation (2.8) defines Yang-Mills gauge field coupled to fermions. Because of the linearity condition $\rho(a+b) = \rho(a) + \rho(b)$ the fermions in our theory turn out to exist in generation. Thanks to $\sum_i a^i b^i = 1$ the inhomogeneous gauge transformation

$${}^g A = gAg^* + gdg^* \quad (2.9)$$

is induced by Eq.(2.6).

One may now ask what happens if fermions are massive. To generalize Eq.(2.2) to this case we define the extra differential to formally produce the mass term using the chiral decomposition of the spinor $\psi = \begin{pmatrix} \psi_L \\ \psi_R \end{pmatrix} \in \mathcal{H}$ by

$$d_\chi \psi = M\psi\chi, \quad M = \begin{pmatrix} 0 & M_1 \\ M_1^\dagger & 0 \end{pmatrix} = M^\dagger \quad (2.10)$$

where M is fermion mass matrix. The extra one-form basis χ combines with the usual ones $\{d\hat{x}^\mu\}_{\mu=0,1,2,3}$ to form 5-dimensional one-form basis:

$$d\hat{x}^\mu \wedge d\hat{x}^\nu = -d\hat{x}^\nu \wedge d\hat{x}^\mu, \quad d\hat{x}^\mu \wedge \chi = -\chi \wedge d\hat{x}^\mu, \quad \chi \wedge \chi = 0 \quad (2.11)$$

Assuming further

$$\langle d\hat{x}^\mu, \chi \rangle = \langle \chi, d\hat{x}^\mu \rangle = 0, \quad \langle \chi, \chi \rangle = -1 \quad (2.12)$$

we obtain the Dirac lagrangian for massive fermion through

$$\mathcal{L}_D = i\langle \tilde{\psi}, \mathbf{d}\psi \rangle, \quad \tilde{\psi} = \gamma_\mu \psi d\hat{x}^\mu + i\psi\chi \quad (2.13)$$

where $\mathbf{d} = d + d_\chi$. As in the massless case Eq.(2.13) is not invariant under Eq.(2.3). But the recipe is now clear. The sum

$$\mathcal{L}_D = i\sum_i \langle a^{i*} \tilde{\psi}, \mathbf{d}(b^i \psi) \rangle, \quad a^i, b^i \in \mathcal{A} \quad (2.14)$$

becomes gauge invariant under Eqs.(2.3) and (2.6). Putting

$$\mathbf{A} = \sum_i a^i \mathbf{d}b^i = A_\mu d\hat{x}^\mu + \Phi\chi \quad (2.15)$$

we rewrite Eq.(2.14) as

$$\mathcal{L}_D = i\langle \tilde{\psi}, (\mathbf{d} + \mathbf{A})\psi \rangle \quad (2.16)$$

Consequently, Eq.(2.14) introduces gauge-invariant coupling of scalars, Φ , χ being assumed to be Lorentz scalar, to fermions.

The gauge transformation property of $\mathbf{A}$ under Eq.(2.6) determines that of Φ in addition to Eq.(2.9):

$${}^g \Phi \chi = g\Phi g^* \chi + gd_\chi g^* \quad (2.17)$$

where use has been made of the relation $\sum_i a^i b^i = 1$. Unless the inhomogeneous term in Eq.(2.17) vanishes Φ is not a physical field.

In writing Eqs.(2.14), (2.15) and (2.16) we have implicitly assumed the Leibniz rule

$$d_\chi(f\psi) = (d_\chi f)\psi + f(d_\chi \psi), \quad f \in \mathcal{A}, \quad \chi\psi = \psi\chi \quad (2.18)$$

Note that with the chiral decomposition of the spinor we must write elements of $\mathcal{A}$ as 2×2 matrix in block form like $f = \begin{pmatrix} f_1 & g_1 \\ g_2 & f_2 \end{pmatrix} \in \mathcal{A}$. It follows that^{*)}

$$d_\chi f = (Mf - fM)\chi, \quad f \in \mathcal{A}, \quad d_\chi M = 0 \quad (2.19)$$

which in turn implies the following Leibniz rule

$$d_\chi(fg) = (d_\chi f)g + f(d_\chi g), \quad f, g \in \mathcal{A}, \quad \chi g = g\chi \quad (2.20)$$

The extended operator $\mathbf{d}$ is nilpotent

$$\mathbf{d}^2 = 0 \quad (2.21)$$

since $d_\chi \chi = 0$ ¹²⁾ and $d^2 = d_\chi^2 = dd_\chi + d_\chi d = 0$. We need to prove the last equality:

$$\begin{aligned} (dd_\chi + d_\chi d)f &= d(Mf - fM)\chi + d_\chi df = (M\partial_\mu f - \partial_\mu fM)(d\hat{x}^\mu \wedge \chi + \chi \wedge d\hat{x}^\mu) = 0 \\ (dd_\chi + d_\chi d)\psi &= d(M\psi)\chi + d_\chi(d\psi) = M\partial_\mu \psi(d\hat{x}^\mu \wedge \chi + \chi \wedge d\hat{x}^\mu) = 0 \end{aligned} \quad (2.22)$$

According to Eq.(2.19) the inhomogeneous term in the right-hand side of Eq.(2.17) vanishes if the mass term is gauge invariant, $[M, g] = 0$. Then Φ is a physical field irrelevant to generation of fermion masses. As an example one may remark that, in this case, there is no need to make the sum (2.14), but

$$\mathcal{L}_D = i \sum_i \langle a^{i*} \tilde{\psi}, d(b^i \psi) \rangle + i \langle \tilde{\psi}, d_\chi \psi \rangle, \quad a^i, b^i \in \mathcal{A} \quad (2.23)$$

is already gauge invariant. Hence we are led back to simply add gauge invariant mass term to Eq.(2.7), leading to no scalars coupled to fermions.

The decomposition (2.15) of the generalized one-form $\mathbf{A}$ into gauge A_μ and shifted Higgs Φ fields corresponds to Connes' prescription of NCG to unify gauge and Higgs fields, where the mass term in the Dirac operator yields the shifted Higgs field. Likewise, our d_χ gives rise to Φ .

We now go on to define the field strength corresponding to $\mathbf{A}$. It will yield the bosonic lagrangian involving solely the bosons coupled to fermions. We employ the definition by Clifford product as proposed in Ref.16).

$$\begin{aligned} \mathbf{G} &= \mathbf{d} \vee \mathbf{A} + \mathbf{A} \vee \mathbf{A} = \mathbf{F} + \mathbf{F}_0 \\ \mathbf{F} &= \mathbf{d} \wedge \mathbf{A} + \mathbf{A} \wedge \mathbf{A} \equiv d\mathbf{A} + \mathbf{A}^2, \quad d\mathbf{A} = \sum_i da^i \wedge db^i \\ \mathbf{F}_0 &= \langle \mathbf{d}, \mathbf{A} \rangle + \langle \mathbf{A}, \mathbf{A} \rangle, \quad \langle \mathbf{d}, \mathbf{A} \rangle = \sum_i \langle da^i, db^i \rangle \end{aligned} \quad (2.24)$$

The field strength $\mathbf{G}$ is inhomogeneous in the rank of differential forms. In other words, it is given by the sum of two-form $\mathbf{F}$ and zero-form $\mathbf{F}_0$ ^{**)} . Both $\mathbf{F}$ and $\mathbf{F}_0$ are gauge covariant so is

^{*)} In the previous papers^{12), 13), 16)} we derived or assumed $d_\chi M \neq 0$. The present modification seems to be more natural.

^{**)} Similarly, Connes' field strength consists of three terms, $\theta = \theta_0 + \theta_1 + \theta_2$ corresponding to two-form $\theta_0 \propto [\gamma^\mu, \gamma^\nu]$, three form $\theta_1 \propto \gamma^\mu \gamma_5$ and zero-form θ_2 containing no Dirac matrices, γ^μ being regarded as one-form basis.

$\mathbf{G}^*$). Explicitly we have

$$\begin{aligned} \mathbf{F} &= F + DH \wedge \chi, \quad F = dA + A^2, \quad DH = dH + AH - HA, \quad H = \Phi + M \\ \mathbf{F}_0 &= X - (H^2 - M^2 - Y), \quad X = -\sum_i a^i \partial^2 b^i + \partial^\mu A_\mu + A^\mu A_\mu = \begin{pmatrix} X_1 & 0 \\ 0 & X_2 \end{pmatrix} \\ A_\mu &= \sum_i a^i \partial_\mu b^i, \quad \Phi = \sum_i a^i [M, b^i], \quad Y = \sum_i a^i [M^2, b^i] = \begin{pmatrix} Y_1 & 0 \\ 0 & Y_2 \end{pmatrix} \end{aligned} \quad (2.25)$$

The system is governed by the lagrangian

$$\mathcal{L} = \mathcal{L}_D + \mathcal{L}_B \quad (2.26)$$

where the fermionic sector is given by Eq.(2.16), while the bosonic sector is obtained by

$$\mathcal{L}_B = -\frac{1}{4} \langle\langle z^2 \mathbf{G}, \mathbf{G} \rangle\rangle = -\frac{1}{4} \langle\langle z^2 \mathbf{F}, \mathbf{F} \rangle\rangle - V, \quad V = \frac{1}{4} \langle\langle z^2 \mathbf{F}_0, \mathbf{F}_0 \rangle\rangle \quad (2.27)$$

where z^2 is a positive matrix commuting with the gauge group and we have used the fact that two-form and zero-form are orthogonal.**)

The lagrangian depends on the model. In the next section we shall discuss toy models.

§3. Toy models

First of all we shall consider $SU(2)$ gauge theory. As is well known there exists an infinite tower of unitary irreps of $SU(2)$ with dimensions $2j+1, j=0, \frac{1}{2}, 1, \dots$. $SU(2)$ gauge theory is defined for the spinor belonging to any one of them.

Now local $SU(2)$ is defined by the unitaries of the algebra $\mathcal{A} = C^\infty(M_4) \otimes \mathbf{H}$. Real quaternion $\mathbf{H}$ has only one irrep

$$\mathbf{H} \ni a \mapsto \begin{pmatrix} \alpha & \beta \\ -\beta^* & \alpha^* \end{pmatrix}, \quad \alpha, \beta \in \mathbf{C} \quad (3.1)$$

Hence one can write $SU(2) = \{g \in \mathbf{H}; gg^* = g^*g = 1, g^* = \bar{g}\}$, where $\bar{g}$ is quaternion conjugation. Using the irrep (3.1) $\mathbf{H}$ is represented by

$$\mathbf{H} \ni a \mapsto \rho(a) = \text{diag}(a, \dots, a^*, \dots), \quad a^* = \text{c.c. of } a \quad (3.2)$$

which acts on a repetition of fundamental reps of $SU(2)$. Consequently, fermions exist in generation and must be non-chiral so that M commutes with $SU(2)$ and $\Phi = \sum_i a^i [M, b^i] = 0$. Thus

*) The fact that both $\mathbf{F}$ and $\mathbf{F}_0$ are separately gauge covariant implies that the most general field strength in this scheme contains an arbitrary real parameter κ : $\mathbf{G} = \mathbf{F} + \kappa \mathbf{F}_0$. Then the arbitrary parameter κ^2 would appear in V . For simplicity in this paper we put $\kappa = 1$ so that the field strength is given by the Clifford product.

**) In evaluating Eq.(2.27) we use the inner product

$$\langle d\hat{x}^\mu \wedge d\hat{x}^\nu, d\hat{x}^\rho \wedge d\hat{x}^\sigma \rangle = g^{\mu\rho} g^{\nu\sigma} - g^{\mu\sigma} g^{\nu\rho}, \quad \langle d\hat{x}^\mu \wedge \chi, d\hat{x}^\nu \wedge \chi \rangle = -g^{\mu\nu}$$

$H = M$ is constant, commuting with A , hence $DH = 0$. We then conclude that $\mathcal{L}_B$ contains only Yang-Mills term because $Y = 0$ and V vanishes due to the equation of motion. That is, our algebraic recipe leads to $SU(2)$ gauge theory without scalars, where fermion doublets exist in generation.

Higgs mechanism can be incorporated into our scheme by enlarging the algebra to $\mathcal{A} = C^\infty(M_4) \otimes (\mathbf{H} \oplus \mathbf{C})$. The unitary group is $\text{Map}(M_4, SU(2) \times U(1)) = \mathcal{U}(C^\infty(M_4) \otimes (\mathbf{H} \oplus \mathbf{C}))$. Writing

$$\mathbf{A} = \sum_i \rho(a_1^i, b_1^i) d\rho(a_2^i, b_2^i) \quad (3.3)$$

for Eq.(2.15) we assume the following rep of $\mathcal{A}$ on the spinor with chiral components

$$\rho(a, b) = \begin{pmatrix} a \otimes 1_{N_g} & 0 \\ 0 & b \otimes 1_{N_g} \end{pmatrix}, \quad a \in \mathbf{H}, \quad b \in \mathbf{C} \quad (3.4)$$

so that left-handed and right-handed spinors belong to doublet and singlet, respectively^{*)}. Except for generation indices ψ_L has two-components like $\begin{pmatrix} \nu_L \\ e_L \end{pmatrix}$, while ψ_R has only one component like e_R . Here and hereafter we omit infinite-dimensional part $C^\infty(M_4)$ of the algebra and remember that, say, a in Eq.(3.4) is $\mathbf{H}$ -valued local functions represented by Eq.(3.1). Choosing the mass matrix as

$$M = \begin{pmatrix} 0 & M_1 \\ M_1^\dagger & 0 \end{pmatrix}, \quad M_1 = \begin{pmatrix} 0 \\ m \end{pmatrix} \quad (3.5)$$

where m is $N_g \times N_g$ mass matrix, we find from the definition of Φ in Eq.(2.25)

$$\Phi = \begin{pmatrix} 0 & \varphi M_1 \\ M_1^\dagger \varphi^\dagger & 0 \end{pmatrix}, \quad \varphi = \begin{pmatrix} \varphi_0^* & \varphi_+ \\ -\varphi_- & \varphi_0 \end{pmatrix} \quad (3.6)$$

Note that φ is $\mathbf{H}$ -valued scalar function. Also

$$H = \Phi + M = \begin{pmatrix} 0 & h M_1 \\ M_1^\dagger h^\dagger & 0 \end{pmatrix} \quad (3.7)$$

$$h = \varphi + 1_2 = \begin{pmatrix} \phi_0^* & \phi_+ \\ -\phi_- & \phi_0 \end{pmatrix}, \quad \phi_0 = \varphi_0 + 1, \quad \phi_\pm = \varphi_\pm$$

It can be shown that Eq.(2.16) leads to

$$\begin{aligned} \mathcal{L}_D &= \bar{\psi}_L i\gamma^\mu (\partial_\mu - \frac{ig_2}{2} A_\mu^{(2)}) \psi_L + \bar{\psi}_R i\gamma^\mu (\partial_\mu - \frac{ig_1}{2} A_\mu^{(1)}) \psi_R \\ &\quad - \bar{\psi}_L m \phi \psi_R - \bar{\psi}_R \phi^\dagger m^\dagger \psi_L \end{aligned} \quad (3.8)$$

where

$$\phi = \begin{pmatrix} \phi_+ \\ \phi_0 \end{pmatrix}, \quad \langle \phi \rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad (3.9)$$

^{*)} There is no reason to assume the same N_g in the doublet and singlet sectors. This assumption is only for later convenience. The $N \times N$ unit matrix is denoted by 1_N .

is the normalized Higgs field and we put

$$A = \sum_i \rho(a_1^i, b_1^i) d\rho(a_2^i, b_2^i) = \begin{pmatrix} -\frac{ig_2}{2} A_\mu^{(2)} d\hat{x}^\mu \otimes 1_{N_g} & 0 \\ 0 & -\frac{ig_1}{2} A_\mu^{(1)} d\hat{x}^\mu \otimes 1_{N_g} \end{pmatrix} \quad (3.10)$$

$A_\mu^{(2)}$ is $SU(2)$ gauge field, while $A_\mu^{(1)}$ $U(1)$ gauge field, both hermitian, g_2 and g_1 being the corresponding gauge coupling constants, respectively.

Let us now evaluate the bosonic lagrangian (2.27) using the formulae (2.25). Putting

$$z^2 = \frac{4}{N_g} \begin{pmatrix} g_2^{-2} 1_2 \otimes 1_{N_g} & 0 \\ 0 & 2g_1^{-2} \otimes 1_{N_g} \end{pmatrix} \quad (3.11)$$

we have for the two-form piece

$$-\frac{1}{4} \langle\langle z^2 \mathbf{F}, \mathbf{F} \rangle\rangle = -\frac{1}{8} \text{tr} F_{\mu\nu}^{(2)} F^{(2)\mu\nu} - \frac{1}{4} F_{\mu\nu}^{(1)} F^{(1)\mu\nu} + L (D^\mu \phi)^\dagger (D_\mu \phi) \quad (3.12)$$

where $F_{\mu\nu}^{(i)}$ are field strength of the gauge fields $A_\mu^{(i)}$, $i = 1, 2$, $D_\mu \phi = (\partial_\mu - \frac{ig_2}{2} A_\mu^{(2)} + \frac{ig_1}{2} A_\mu^{(1)}) \phi$ and we define

$$L = \frac{2}{N_g} \left(\frac{1}{g_2^2} + \frac{2}{g_1^2} \right) \text{tr}_g (m^\dagger m) \quad (3.13)$$

with tr_g denoting the trace in the generation space. Next we compute the zero form piece. Since $Y_2 = 0$ in Eq.(2.25), X_1, X_2 and Y_1 are auxiliary fields to be eliminated from the lagrangian using the equation of motion, we get

$$V = K(\phi^\dagger \phi - 1)^2, \quad K = \frac{2}{N_g g_1^2} (\text{tr}_g (M_1^\dagger M_1)^2 - \frac{1}{N_g} (\text{tr}_g M_1^\dagger M_1)^2) \quad (3.14)$$

Finally, we have

$$\mathcal{L}_B = -\frac{1}{8} \text{tr} F_{\mu\nu}^{(2)} F^{(2)\mu\nu} - \frac{1}{4} F_{\mu\nu}^{(1)} F^{(1)\mu\nu} + (D^\mu \phi)^\dagger (D_\mu \phi) - \frac{\lambda}{4} (\phi^\dagger \phi - \frac{v^2}{2})^2 \quad (3.15)$$

where

$$\lambda = \frac{4K}{L^2}, \quad v^2 = 2L \quad (3.16)$$

For λ should be positive, the theory is consistently defined only if $N_g > 1$.

The rescaling $\phi \rightarrow \frac{1}{\sqrt{L}} \phi = \frac{\sqrt{2}}{v} \phi$ to obtain above $\mathcal{L}_B$ renders the mass term in Eq.(3.8) multiplied by $\frac{\sqrt{2}}{v}$.

There is another rep of the same algebra other than (3.4):

$$\rho(a, b) = \begin{pmatrix} a \otimes 1_{N_g} & 0 \\ 0 & B \otimes 1_{N_g} \end{pmatrix}, \quad B = \begin{pmatrix} b & 0 \\ 0 & b^* \end{pmatrix} \quad (3.17)$$

which is suitable for 'massive' neutrino. This rep acts on doublet like $\psi_L = \begin{pmatrix} \nu \\ e \end{pmatrix}_L$ and singlets like $\psi_R = \begin{pmatrix} \nu_R \\ e_R \end{pmatrix}$ with generation indices omitted. The mass matrix is given by

$$M_1 = \begin{pmatrix} m_1 & 0 \\ 0 & m_2 \end{pmatrix}, \quad m_{1,2} : N_g \times N_g \quad (3.18)$$

The Higgs field is given by Eqs.(3.6) and (3.7) with M_1 being of the form (3.18). Putting

$$\begin{aligned} A &= \sum_i \rho(a_1^i, b_1^i) d\rho(a_2^i, b_2^i) \\ &= \begin{pmatrix} -\frac{ig_2}{2} A_\mu^{(2)} d\hat{x}^\mu \otimes 1_{N_g} & 0 \\ 0 & -\frac{ig_1}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} A_\mu^{(1)} d\hat{x}^\mu \otimes 1_{N_g} \end{pmatrix} \end{aligned} \quad (3.19)$$

we find

$$\begin{aligned} \mathcal{L}_D &= \bar{\psi}_L i\gamma^\mu (\partial_\mu - \frac{ig_2}{2} A_\mu^{(2)}) \psi_L + \bar{\psi}_R i\gamma^\mu (\partial_\mu - \frac{ig_1}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} A_\mu^{(1)}) \psi_R \\ &\quad - \bar{\psi}_L (m_1 \tilde{\phi}, m_2 \phi) \psi_R - \bar{\psi}_R \begin{pmatrix} \tilde{\phi}^\dagger m_1^\dagger \\ \phi^\dagger m_2 \end{pmatrix} \psi_L, \quad \tilde{\phi} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \phi^* \end{aligned} \quad (3.20)$$

The bosonic lagrangian takes the same form as (3.15) with L of Eq.(3.16) given by

$$L = \frac{2}{N_g} \left(\frac{1}{g_2^2} + \frac{1}{g_1^2} \right) \text{tr}_g (M_1^\dagger M_1) \quad (3.21)$$

provided we set

$$z^2 = \frac{4}{N_g} \begin{pmatrix} g_2^{-2} 1_2 \otimes 1_{N_g} & 0 \\ 0 & g_1^{-2} 1_2 \otimes 1_{N_g} \end{pmatrix} \quad (3.22)$$

Next consider the case $N_g=1$ and let $\psi^a (a=1,2,3)$ belong to the adjoint rep of $SU(2)$ with the matrix field

$$\psi = \tau_a \psi^a \quad (3.23)$$

where $\tau = (\tau_1, \tau_2, \tau_3)$ is the Pauli matrix. Choosing the algebra $\mathcal{A} = C^\infty(M_4) \otimes M_2(\mathbf{C})$ whose unitary group is $\text{Map}(M_4, SU(2) \times U(1)) = \mathcal{U}(C^\infty(M_4) \otimes M_2(\mathbf{C}))$, although $U(1)$ factor is automatically eliminated in the reconstruction below, we represent it by

$$\rho(a) = \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}, \quad a \in M_2(\mathbf{C}) \quad (3.24)$$

Action by $\mathbf{d} + \mathbf{A}$ on ψ is defined by the commutator

$$\begin{aligned} (\mathbf{d} + \mathbf{A})\psi &\equiv [\mathbf{d} + \mathbf{A}, \psi] = [d + A, \psi] + [d_\chi + \Phi\chi, \psi] \\ [d + A, \psi] &\equiv d\psi + \begin{pmatrix} [A_1, \psi_L] \\ [A_2, \psi_R] \end{pmatrix}, A = \begin{pmatrix} A_1 & 0 \\ 0 & A_2 \end{pmatrix} \\ [d_\chi, \psi] &= [M, \psi] \equiv \begin{pmatrix} [M_1, \psi_R] \\ [M_1^\dagger, \psi_L] \end{pmatrix}, [\Phi, \psi] \equiv \begin{pmatrix} [\varphi, \psi_R] \\ [\varphi^\dagger, \psi_L] \end{pmatrix} \end{aligned} \quad (3.25)$$

where

$$\Phi = \sum_i \rho(a^i)[M, \rho(b^i)] = \begin{pmatrix} 0 & \varphi \\ \varphi^\dagger & 0 \end{pmatrix}, \quad a^i, b^i \in \mathcal{A} \quad (3.26)$$

By assumption $A_1 = A_2$ and we put $A_{1\mu} = -igA_\mu$ where A_μ is hermitian $SU(2)$ gauge field due to the commutator (3.25). Hence the gauge group is $SU(2)$, not $SU(2) \times U(1)$. Another comment here is that Φ does not take the form (3.6) but has three complex components in view of the commutator (3.25):

$$\varphi = \begin{pmatrix} \varphi_{11} & \varphi_{12} \\ \varphi_{21} & -\varphi_{11} \end{pmatrix} \quad (3.27)$$

$H = \Phi + M$ is given by Eq.(3.27) with $\varphi \rightarrow h = \varphi + M_1$, whence we put

$$-\frac{1}{g}h \equiv \phi = \tau_a \phi^a \equiv S - iP \quad (3.28)$$

It is now easy to evaluate¹⁵⁾ the lagrangian

$$\mathcal{L}_B = -\frac{1}{4g^2} \langle\langle \mathbf{G}, \mathbf{G} \rangle\rangle = -\frac{1}{4g^2} \langle\langle \mathbf{F}, \mathbf{F} \rangle\rangle - V, \quad V = \frac{1}{4g^2} \langle\langle \mathbf{F}_0, \mathbf{F}_0 \rangle\rangle \quad (3.29)$$

where we put $z^2 = \frac{1}{g^2}1_4$ in Eq.(2.27) to take into account supersymmetry. The result is

$$\mathcal{L}_B = -\frac{1}{8} \text{tr} F_{\mu\nu} F^{\mu\nu} + \frac{1}{2} \text{tr} ([D^\mu, \phi])^\dagger [D_\mu, \phi] - \frac{g^2}{8} \text{tr} [\phi, \phi^\dagger]^2 \quad (3.30)$$

where $D_\mu = \partial_\mu - igA_\mu$, provided M_1 is a normal matrix, $M_1 M_1^\dagger = M_1^\dagger M_1$ ^{*)}.

In the fermionic sector we simply find from Eqs.(2.16), (3.25), (3.26) and (3.28)

$$\mathcal{L}_D = \text{tr}(\bar{\psi} i \gamma^\mu [D_\mu, \psi] + g \bar{\psi} [S + i \gamma_5 P, \psi]) \quad (3.31)$$

As is well known the sum of Eqs.(3.30) and (3.31) defines $N = 2$ super Yang-Mills theory²⁰⁾.

§4. Standard model of elementary particles

To reconstruct the standard model of elementary particles in the present scheme we take into account Connes real structure J . To define J as simple as possible we follow Connes to

^{*)} This implies that $Y = 0$ in Eq.(2.25). Hence there must be X term in $\mathbf{F}_0$ to obtain the scalar potential.

double spinors using charge conjugate spinor ψ^c as

$$\Psi = \begin{pmatrix} \psi \\ \psi^c \end{pmatrix}, \quad \psi = \begin{pmatrix} \psi_L \\ \psi_R \end{pmatrix} \quad (4.1)$$

upon which gauge transformation acts like

$${}^g\Psi = U\Psi = U_1 U_2 \Psi, \quad U^\dagger U = U_i^\dagger U_i = 1 \quad (4.2)$$

if

$$U = \begin{pmatrix} u_1 u_2^* & 0 \\ 0 & u_1^* u_2 \end{pmatrix}, \quad u_i^* = \text{c.c. of } u_i \quad (4.3)$$

where

$$U_1 = \begin{pmatrix} u_1 & 0 \\ 0 & u_2 \end{pmatrix}, \quad U_2 = \begin{pmatrix} u_2^* & 0 \\ 0 & u_1^* \end{pmatrix} \quad (4.4)$$

with $u_1^* u_2 = u_2 u_1^*$, or $U_1 U_2 = U_2 U_1$. In the language of Connes mathematics obtaining U_2 from U_1 is accomplished by means of the real structure J , $U_2 = J U_1 J^\dagger$, provided U_1 is a rep of a noncommutative algebra^{*)}. Since Abelian charges cancel among fermions and anti-fermions, $\det U = 1$. This condition is satisfied if

$$\det u_1 u_2^* = 1 \quad (4.5)$$

which is highly nontrivial. We shall refer Eq.(4.5) to as Connes unimodularity condition.

In the standard model we choose the basis using the mass eigenstates

$$\psi = \begin{pmatrix} q_L \\ l_L \\ u_R \\ d_R \\ \nu_{eR} \\ e_R \end{pmatrix}, \quad q_L = \begin{pmatrix} u \\ U_q d \end{pmatrix}_L, \quad l_L = \begin{pmatrix} \nu_e \\ U_l e \end{pmatrix}_L \quad (4.6)$$

where we omit generation indices on leptons and quarks as well as color indices on quarks and similarly for anti-fermions. The $N_g \times N_g$ Kobayashi-Maskawa matrices are denoted by $U_{l,q}$ in the lepton and quark sectors, respectively. Experiment shows $N_g = 3$ up to the present energy but we let N_g be a free parameter. We assume massive neutrinos using the rep (3.17) in the lepton sector.

To recover the standard gauge group $SU(3) \times SU(2)_L \times U(1)_Y$ from $\mathcal{U}(\mathcal{A})$ we make Connes choice^{4), 5)}

$$\mathcal{A} = C^\infty(M_4) \otimes (\mathbf{H} \oplus \mathbf{C} \oplus M_3(\mathbf{C})) \quad (4.7)$$

whose unitaries are $\text{Map}(M_4, U(3) \times SU(2) \times U(1))$. We shall see below that the unimodularity condition (4.5) reduces $U(3) \times SU(2) \times U(1)$ to $SU(3) \times SU(2)_L \times U(1)_Y$ with correct hypercharge assignment.

^{*)} The operator J is an anti-linear isometry in the total Hilbert space $\mathcal{H}$ which satisfies two conditions^{4), 5)}. Then Eq.(4.2) reads ${}^g\Psi = U_1 \Psi U_1^* \equiv U_1 J U_1 J^* \Psi$.

In the basis (4.6) Eqs.(4.2), (4.3) and (4.4) turn out to be given by the unitary restriction of the rep of Eq.(4.7): $u_1 = \rho_w(a, b)$ and $u_2 = \rho_s(b, c)$ for $(a, b, c) \in \mathcal{U}(\mathcal{A})$ where, for general $a \in \mathbf{H}, b \in \mathbf{C}, c \in M_3(\mathbf{C})$, $\mathcal{A}$ is represented on Eq.(4.1) by ^{4), 5), 17), 18), 19)}

$$\begin{aligned} \rho(a, b, c) &= \begin{pmatrix} \rho_w(a, b) & 0 \\ 0 & \rho_s(b, c) \end{pmatrix} \\ \rho_w(a, b) &= \begin{pmatrix} \rho_1(a) & 0 \\ 0 & \rho_2(b) \end{pmatrix}, \quad \rho_s(b, c) = \begin{pmatrix} \rho_3(b, c) & 0 \\ 0 & \rho_4(b, c) \end{pmatrix} \\ \rho_1(a) &= \begin{pmatrix} a \otimes 1_3 \otimes 1_{N_g} & 0 \\ 0 & a \otimes 1_{N_g} \end{pmatrix}, \quad \rho_2(b) = \rho_1(B) \\ \rho_3(b, c) &= \begin{pmatrix} 1_2 \otimes c^* \otimes 1_{N_g} & 0 \\ 0 & b 1_2 \otimes 1_{N_g} \end{pmatrix} = \rho_4(b, c) \end{aligned} \quad (4.8)$$

where B is given by Eq.(3.17). By U_1 flavor acts only in the particle sector, whereas color operates only in the anti-particle sector*).

The well-known fermionic sector of the standard model lagrangian is given using the doubled spinor (4.1) by

$$\mathcal{L}_D = \frac{i}{2} \langle \tilde{\Psi}, (\mathbf{d} + \mathbf{A} + J \mathbf{A} J^\dagger) \Psi \rangle \quad (4.9)$$

where

$$\begin{aligned} d_\chi \Psi &= \mathcal{M} \Psi \chi, \quad \mathcal{M} = \mathcal{M}_1 + J \mathcal{M}_1 J^\dagger, \quad \mathcal{M}_1 = \begin{pmatrix} M & 0 \\ 0 & 0 \end{pmatrix}, \quad M = \begin{pmatrix} 0 & M_1 \\ M_1^\dagger & 0 \end{pmatrix} \\ \mathbf{A} &= \sum_i \rho(a_1^i, b_1^i, c_1^i) \mathbf{d} \rho(a_2^i, b_2^i, c_2^i) = A + \Phi \chi \\ A &= \begin{pmatrix} A_w & 0 \\ 0 & A_s \end{pmatrix}, \quad A_w = \sum_i \rho_w(a_1^i, b_1^i) d \rho_w(a_2^i, b_2^i), \quad A_s = \sum_i \rho_s(b_1^i, c_1^i) d \rho_s(b_2^i, c_2^i) \\ \Phi &= \sum_i \rho(a_1^i, b_1^i, c_1^i) [\mathcal{M}, \rho(a_2^i, b_2^i, c_2^i)] = \begin{pmatrix} \begin{pmatrix} 0 & \rho_1(\varphi) M_1 \\ M_1^\dagger \rho_1^\dagger(\varphi) & 0 \end{pmatrix} & 0 \\ 0 & 0 \end{pmatrix} \\ H &= \Phi + \mathcal{M}_1 = \begin{pmatrix} \begin{pmatrix} 0 & \rho_1(h) M_1 \\ M_1^\dagger \rho_1^\dagger(h) & 0 \end{pmatrix} & 0 \\ 0 & 0 \end{pmatrix} \end{aligned} \quad (4.10)$$

with

$$J \begin{pmatrix} p & 0 \\ 0 & q \end{pmatrix} J^\dagger = \begin{pmatrix} q^* & 0 \\ 0 & p^* \end{pmatrix} \quad (4.11)$$

^{*)} Before introduction of the real structure, Connes considered ³⁾ the bimodule $\mathcal{A} \oplus \mathcal{B}$, $\mathcal{A} = \mathbf{H} \oplus \mathbf{C}$, $\mathcal{B} = \mathbf{C} \oplus M_3(\mathbf{C})$ with left action by flavor and right action by color. Thanks to the real structure J , the bimodule structure of $\mathcal{H}$ is obtained with single algebra (4.7) just because there exist matter and anti-matter in quantum field theory.

Here φ is defined by Eq.(3.6), h by Eq.(3.7) and $\rho_1(a), a \in \mathbf{H}$ by Eq.(4.8). Note that in the basis (4.6) we have the following mass matrix:

$$\begin{aligned} M_1 &= \begin{pmatrix} M_q \otimes 1_3 & 0 \\ 0 & M_l \end{pmatrix}, \quad M_q = \begin{pmatrix} M_u & 0 \\ 0 & M_d \end{pmatrix}, \quad M_l = \begin{pmatrix} M_\nu & 0 \\ 0 & M_e \end{pmatrix} \\ M_u &= \text{diag}(m_u, m_c, m_t, \dots), \quad M_d = U_q \text{diag}(m_d, m_s, m_b, \dots) \\ M_\nu &= \text{diag}(m_{\nu_e}, m_{\nu_\mu}, m_{\nu_\tau}, \dots), \quad M_e = U_l \text{diag}(m_e, m_\mu, m_\tau, \dots) \end{aligned} \quad (4.12)$$

with obvious notations for fermion masses. KM matrices $U_f, f = l, q$ disappear due to the corresponding factors in Eq.(4.6). Recall that $M_f, f = q, l$ are assumed to be $2N_g \times 2N_g$ matrices.

In Eq.(4.9) we have two generalized gauge potentials, $\mathbf{A}$ and $J\mathbf{A}J^\dagger$. They correspond to two factors, U_1 and U_2 in Eq.(4.2). In fact, we see from the definition (4.10) and (4.11)

$$\begin{aligned} {}^g\mathbf{A} &= U_1 \mathbf{A} U_1^\dagger + U_1 dU_1^\dagger, \quad U_2 \mathbf{A} U_2^\dagger = \mathbf{A} \\ J^g\mathbf{A}J^\dagger &= U_2 (J\mathbf{A}J^\dagger) U_2^\dagger + U_2 dU_2^\dagger, \quad U_1 (J\mathbf{A}J^\dagger) U_1^\dagger = J\mathbf{A}J^\dagger \\ {}^g\mathbf{A} + J^g\mathbf{A}J^\dagger &= U(\mathbf{A} + J\mathbf{A}J^\dagger)U^\dagger + U dU^\dagger \end{aligned} \quad (4.13)$$

where we have used the relation $U_1(U_2 dU_2^\dagger)U_1^\dagger = U_2 dU_2^\dagger$. Hence the fermionic lagrangian (4.9) is gauge invariant under (4.2). What remains to be proved is that the unimodularity condition (4.5) determines the correct hypercharge assignment ^{2), 3), 4), 10)}. To see what is happening in the theory, we set

$$\begin{aligned} A_w &= \begin{pmatrix} A_w^L & 0 \\ 0 & A_w^R \end{pmatrix}, \quad A_s = \begin{pmatrix} A_s^L & 0 \\ 0 & A_s^R \end{pmatrix} \\ A_w^L &= \begin{pmatrix} W \otimes 1_3 \otimes 1_{N_g} & 0 \\ 0 & W \otimes 1_{N_g} \end{pmatrix} \\ A_w^R &= W \rightarrow \begin{pmatrix} B^* & 0 \\ 0 & B \end{pmatrix} \text{ in } A_w^L \\ A_s^L &= \begin{pmatrix} 1_2 \otimes G^* \otimes 1_{N_g} & 0 \\ 0 & \begin{pmatrix} B^* & 0 \\ 0 & B^* \end{pmatrix} \otimes 1_{N_g} \end{pmatrix} = A_s^R \end{aligned} \quad (4.14)$$

where

$$W = \sum_i a_1^i da_2^i, \quad B^* = \sum_i b_1^i db_2^i, \quad G = \sum_i c_1^i dc_2^i \quad (4.15)$$

are flavor, Abelian and "color" gauge fields. The last still contains Abelian part

$$G = G' + \frac{1}{3} 1_3 \text{tr} G, \quad \text{tr} G' = 0 \quad (4.16)$$

Now the unimodularity condition (4.5) means that $\text{tr}(A_w + A_s^*) = 0$. Noting that W and G' are traceless and gauge fields are chosen as antihermitian, we get at

$$B + \text{tr} G = 0 \quad (4.17)$$

indicating that there is only one Abelian gauge field, which we take to be $U(1)_Y$ gauge field B . If B couples to lepton doublet with strength proportional to $-y$, we are led to the following hypercharge assignment

$$\begin{aligned} Y_L^l &= \begin{pmatrix} -y & 0 \\ 0 & -y \end{pmatrix}, & Y_L^q &= \begin{pmatrix} \frac{1}{3}y & 0 \\ 0 & \frac{1}{3}y \end{pmatrix} \otimes 1_3 \\ Y_R^l &= \begin{pmatrix} 0 & 0 \\ 0 & -2y \end{pmatrix}, & Y_R^q &= \begin{pmatrix} y + \frac{1}{3}y & 0 \\ 0 & -y + \frac{1}{3}y \end{pmatrix} \otimes 1_3 \end{aligned} \quad (4.18)$$

By rescaling

$$B \rightarrow -\frac{ig'}{2}B, \quad W \rightarrow -\frac{ig}{2}W, \quad G' \rightarrow -\frac{ig_s}{2}G' \quad (4.19)$$

and putting $y = +1$ we are able to recover the fermionic lagrangian of the standard model from Eq.(4.9) with normalized Higgs field.

Next we turn to the bosonic sector where we make use of only $\mathbf{A}$ in Eq.(4.10) to define the field strength $\mathbf{G}$. The formula (2.27) gives

$$\mathcal{L}_B = \mathcal{L}_{YM} + \mathcal{L}_H^{K.E.} - V \quad (4.20)$$

where, putting

$$z^2 = \begin{pmatrix} z_p^2 & 0 \\ 0 & z_a^2 \end{pmatrix}, \quad \left\{ \begin{aligned} z_p^2 &= \begin{pmatrix} z_1^2 1_{6N_g} & & & 0 \\ & z_1'^2 1_{2N_g} & & \\ & & z_2^2 1_{6N_g} & \\ & & & z_2'^2 1_{2N_g} \end{pmatrix} \\ z_a^2 &= \begin{pmatrix} & & & 0 \\ z_3^2 1_{6N_g} & & & \\ & z_3'^2 1_{2N_g} & & \\ 0 & & z_4^2 1_{6N_g} & \\ & & & z_4'^2 1_{2N_g} \end{pmatrix} \end{aligned} \right\} \quad (4.21)$$

with the conditions

$$\begin{aligned} (3z_1^2 + z_1'^2) \frac{g^2}{4} N_g &= 1 \\ (3z_2^2 + z_2'^2) \frac{g'^2}{2} N_g + (z_3^2 + z_4^2 + 3(z_3'^2 + z_4'^2)) \frac{g'^2}{6} N_g &= 2 \\ (z_3^2 + z_4^2) \frac{g_s^2}{2} N_g &= 1 \end{aligned} \quad (4.22)$$

Yang-Mills term becomes

$$\mathcal{L}_{YM} = -\frac{1}{8} \text{tr}(W_{\mu\nu} W^{\mu\nu} + G'_{\mu\nu} G'^{\mu\nu}) - \frac{1}{4} B_{\mu\nu} B^{\mu\nu} \quad (4.23)$$

in terms of $SU(2)_L, SU(3), U(1)_Y$ gauge field strengths, $W_{\mu\nu}, G'_{\mu\nu}, B_{\mu\nu}$, respectively, and Higgs kinetic energy term is given by

$$\mathcal{L}_H^{K.E.} = L(D^\mu \phi)^\dagger (D_\mu \phi) \quad (4.24)$$

with

$$L = \frac{1}{4} \text{tr}_g [3(z_1^2 + z_2^2)(M_u M_u^\dagger + M_d M_d^\dagger) + (z_1'^2 + z_2'^2)(M_\nu M_\nu^\dagger + M_e M_e^\dagger)] \quad (4.25)$$

Note that ϕ in (4.24) is normalized such that $\langle \phi \rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$. On the other hand, the potential V still contains auxiliary fields to be eliminated to take the following form

$$V = K(\phi^\dagger \phi - 1)^2 \quad (4.26)$$

with

$$K = \frac{1}{4} \left(3z_2^2 \sum_{q=u,d} [\text{tr}_g(M_q M_q^\dagger)^2 - \frac{1}{N_g} (\text{tr} M_q M_q^\dagger)^2] + z_2'^2 \sum_{l=\nu,e} [\text{tr}_g(M_l M_l^\dagger)^2 - \frac{1}{N_g} (\text{tr} M_l M_l^\dagger)^2] \right) \quad (4.27)$$

The final form of the bosonic lagrangian after rescaling $\phi \rightarrow \frac{1}{\sqrt{L}}\phi$ reads

$$\mathcal{L}_B = -\frac{1}{8} \text{tr}(W^2 + G'^2) - \frac{1}{4} B^2 + (D^\mu \phi)^\dagger (D_\mu \phi) - \frac{\lambda}{4} (\phi^\dagger \phi - \frac{v^2}{2})^2 \quad (4.28)$$

$W^2 = W_{\mu\nu} W^{\mu\nu}$ and so on in Yang-Mills sector, with

$$\lambda = \frac{4K}{L^2}, \quad v^2 = 2L \quad (4.29)$$

The present reconstruction of the standard model requires $N_g > 1$ in order to have $\lambda > 0$. That the restriction $N_g > 1$ is obtained in NCG was first observed by Connes-Lott¹⁾. It should be noted that X term in Eq.(2.25) is necessary to deduce this conclusion, though both X and the rest in Eq.(2.25) are separately gauge covariant. See the footnote on p.12 as regards $N = 2$ super Yang-Mills theory. Therefore, our conclusion $N_g > 1$ depends on the assumption that the present formalism works also for $N = 2$ super Yang-Mills theory. For instance, we never come across the same conclusion if we modify $\mathbf{G} \rightarrow \mathbf{G}' = \mathbf{F} + \mathbf{F}_0'$, where $\mathbf{F}_0' = \sum_i \langle d_\chi a^i, d_\chi b^i \rangle + \langle \Phi_\chi, \Phi_\chi \rangle$.

However, this modification spoils the previous success in reconstructing $N = 2$ super Yang-Mills theory in contrast with NCG consideration¹⁴⁾. The detailed derivation of Eqs.(4.25) and (4.27) will be postponed to the Appendix.

§5. Discussions

Using the relation $m_W^2 = \frac{1}{4} g^2 v^2 = \frac{1}{2} g^2 L$ (see Eq.(4.29)) and estimating L by top quark mass dominance for $N_g = 3$ we find

$$m_W = m_t \sqrt{\frac{z_1^2 + z_2^2}{2(3z_1^2 + z_1'^2)}}$$

If $z_1^2 = z_2^2$, it follows¹⁷⁾ that

$$m_t \geq \sqrt{3}m_W \simeq 139 \text{ GeV}$$

Hence, we are not doing so bad business, although, in fact, we get no constraints among the parameters due to the most general form of the matrix z^2 in Eq.(4.23).

Nevertheless, one may assume that NCG-based bosonic lagrangian with z^2 proportional to the unit matrix 1_{32N_g} should become exact at some large energy scale Λ for unknown reason. Then at Λ we get GUT relations $g_s^2 = g^2 = \frac{5}{3}g'^2$ and the top mass dominance even at such large energy scale would lead to the mass relations $m_t = 2m_W$ and $m_H = \sqrt{2}m_t$ ^{1), 11)*}). We shall not pursue quantitative analysis^{17), 21)} in terms of renormalization group equations of these or similar constraints in this paper.

À la Connes we are led to the rep (4.8) by experiment if $N_g = 3$. We do not yet understand, however, what determines the rep (4.8)**), including the precise value of N_g except for the theoretical restriction $N_g > 1$. Nonetheless the concept of generation is introduced into the theory in a very natural way. And this is accompanied by promoting Higgs field to a kind of gauge field as in more rigorous treatment of Connes NCG. In this sense the present approach, or more general approach of NCG offers a new insight into the Yang-Mills-Higgs theory. It should also be mentioned^{17), 21)} that Higgs mass is estimated to be below or around the weak scale $v = 247$ GeV upon quantum corrections.

Some authors¹⁸⁾ pointed out that the fact that the correct hypercharge assignment (4.18) comes only from Connes unimodularity condition (4.5) indicates that NCG-based standard model should be unified with gravity at large energy scale²²⁾ because Eq.(4.15) or (4.18) assures the absence of graviton-graviton- $U(1)$ anomaly, $\text{tr}Y=0$. Our next task is to construct standard-model-gravity-coupled model based on the present formalism.

*) If we introduce the parameter κ^2 in the expression for V of Eq.(2.27), the mass relation becomes $m_H = \sqrt{2}\kappa m_t$, which makes it unable to predict Higgs mass by means of renormalization group equations.

**) We may argue that the rep (4.8) is determined by the requirements that 1) left-handed fermions be doublets, 2) right-handed fermions singlets, 3) color nonchiral, 4) u_R and d_R have different Abelian charges, 5) ν_R be neutral under $U(3) \times SU(2) \times U(1)$ in each generation. It should be noted, however, that these requirements are all guided by experiment.

Appendix A

— *Appendix* —

Let us first derive Higgs kinetic energy term (4.24). By definition we have

$$\mathcal{L}_H^{K.E.} = \frac{1}{4} \text{tr} z^2 (D_\mu H)^\dagger (D^\mu H) \quad (\text{A.1})$$

where

$$D_\mu H = \begin{pmatrix} \begin{pmatrix} 0 & D_\mu \rho_1(h) M_1 \\ M_1^\dagger (D_\mu \rho_1(h))^\dagger & 0 \end{pmatrix} & \begin{pmatrix} 0 \\ 0 \end{pmatrix} \end{pmatrix} \quad (\text{A.2})$$

with

$$D_\mu \rho_1(h) = (D_\mu \tilde{\phi}, D_\mu \phi) \otimes 1_{4N_g}, \quad \begin{cases} D_\mu \phi = (\partial_\mu - \frac{ig}{2} W_\mu - \frac{ig'}{2} 1_2 B_\mu) \phi \\ D_\mu \tilde{\phi} = (\partial_\mu - \frac{ig}{2} W_\mu + \frac{ig'}{2} 1_2 B_\mu) \tilde{\phi} \end{cases} \quad (\text{A.3})$$

We then compute (A.1) using the relation

$$\begin{aligned} \text{tr}(D^\mu \tilde{\phi}, D^\mu \phi) (M_1 M_1^\dagger) \begin{pmatrix} (D_\mu \tilde{\phi})^\dagger \\ (D_\mu \phi)^\dagger \end{pmatrix} &= \text{tr} M_1^\dagger \begin{pmatrix} (D_\mu \tilde{\phi})^\dagger \\ (D_\mu \phi)^\dagger \end{pmatrix} (D^\mu \tilde{\phi}, D^\mu \phi) M_1 \\ &= \text{tr}_g [3(M_u M_u^\dagger + M_d M_d^\dagger) + (M_\nu M_\nu^\dagger + M_e M_e^\dagger)] (D^\mu \phi)^\dagger (D_\mu \phi) \end{aligned} \quad (\text{A.4})$$

and (4.21) to obtain Eqs.(4.24) and (4.25).

Next we evaluate Eq.(4.27). According to Eq.(2.25) we have

$$\begin{aligned} X &= \begin{pmatrix} X_1 & 0 \\ 0 & X_2 \end{pmatrix} = \begin{pmatrix} \begin{pmatrix} x_1 & 0 \\ 0 & y_1 \end{pmatrix} & 0 \\ 0 & \begin{pmatrix} x_2 & 0 \\ 0 & y_2 \end{pmatrix} \end{pmatrix} \\ H^2 &= \begin{pmatrix} \begin{pmatrix} \Theta & 0 \\ 0 & \Theta' \end{pmatrix} & 0 \\ 0 & 0 \end{pmatrix} \\ M^2 &= \begin{pmatrix} \begin{pmatrix} M_1 M_1^\dagger & 0 \\ 0 & M_1^\dagger M_1 \end{pmatrix} & 0 \\ 0 & 0 \end{pmatrix} \\ Y &= \begin{pmatrix} \begin{pmatrix} Y_1 & 0 \\ 0 & Y_2 \end{pmatrix} & 0 \\ 0 & 0 \end{pmatrix} \end{aligned} \quad (\text{A.5})$$

with

$$\begin{aligned} \Theta &= \rho_1(h) M_1 M_1^\dagger \rho_1^\dagger(h), \quad \Theta' = M_1^\dagger \rho_1^\dagger(h) \rho_1(h) M_1 \\ Y_1 &= \sum_i \rho_1(a_1^i, b_1^i) [M_1 M_1^\dagger, \rho_1(a_2^i, b_2^i)], \quad Y_2 = 0 \end{aligned} \quad (\text{A.6})$$

It is now easy to eliminate the auxiliary fields, $x_1 + Y_1, x_2, y_2$ from the expression for the potential V and determine y_1 in the quark and lepton sectors from the equation of motion:

$$\begin{aligned} y_1^q &= \frac{1}{N_g}(\phi^\dagger \phi - 1) \begin{pmatrix} \text{tr}_g M_u^\dagger M_u & 0 \\ 0 & \text{tr}_g M_d^\dagger M_d \end{pmatrix} \otimes 1_3 \otimes 1_{N_g} \\ y_1^l &= \frac{1}{N_g}(\phi^\dagger \phi - 1) \begin{pmatrix} \text{tr}_g M_\nu^\dagger M_\nu & 0 \\ 0 & \text{tr}_g M_e^\dagger M_e \end{pmatrix} \otimes 1_{N_g} \end{aligned} \quad (\text{A.7})$$

Consequently, we arrive at the final form of the Higgs potential (4.26) with the coefficient (4.27).

References

- [1] A. Connes, p.9 in *The Interface of Mathematics and Particle Physics*, CLARENDON PRESS, OXFORD, 1990.
- [2] A. Connes and J. Lott, Nucl. Phys. Proc. Suppl. **18B**(1990), 29.
- [3] A. Connes, *Noncommutative Geometry*, Academic Press, New York, 1994.
- [4] A. Connes, J. Math. Phys. **36**(1995), 6194.
- [5] A. Connes, Comm. Math. Phys. **182**(1996), 155.
- [6] D. Kastler, Reviews in Math.Phys.**5**(1990), 477.
- [7] J. C. Várilly and J. M. Gracia-Bondia, J.Geom.Phys.**12**(1995), 223;
- [8] B. Iochum and T. Schücker, Comm. Math. Phys. **178**(1996),1.
- [9] R. Coquereaux, G. Esposito-Farese and J. Frölich, Nucl.Phys.**B353**(1991), 689.
- [10] R. Coquereaux, R. Häßling, N. A. Papadopoulos and F. Scheck, Int.J.Mod.Phys. **A7**(1992), 2809.
- [11] K. Morita, Prog.Theor.Phys.**90**(1993), 219.
- [12] S. Naka and E. Umezawa, Prog. Theor. Phys. **92**(1994), 189.
- [13] A. H. Chamseddine, G. Felder and J. Frölich, Phys. Letters **B296**(1992), 109;
- [14] Nucl. Phys. **B395**(1993), 672; Comm. Math. Phys. **155**(1993), 205;
- [15] A. H. Chamseddine and J. Frölich, Phys. Letters **B314**(1993), 308; Phys. Rev. **D50**(1994), 2893;
- [16] A. H. Chamseddine, Phys. Letters **B373**(1996), 61.
- [17] T. Schücker and J.-M. Zylinski, J.Geom.Phys.**16**(1995), 207.
- [18] A. Sitarz, Phys. Lett. **B308**(1993), 311;
- [19] H.-G. Ding, H.-Y. Guo, J.-M. Li and K. Wu, Z.Phys.**C64**(1994), 521;
- [20] B. Chen, T. Saito, H.-B. Teng, K. Uehara and K. Wu, Prog. Theor. Phys. **95**(1996), 1173.
- [21] E. Alvarez, J. M. Gracia-Bondía and C. P. Martin, Phys.Lett.**364**(1995), 33.
- [22] I. S. Sogami, Prog. Theor. Phys. **94**(1995), 117;**95**(1996), 637.
- [23] K. Morita and Y. Okumura, Prog. Theor. Phys. **91**(1994), 959;Phys. Rev. **D50**(1994),1016.
- [24] Y. Okumura, Phys.Rev.**D50**(1994), 1026;Prog. Theor. Phys. **96**(1996), 1021.
- [25] A. H. Chamseddine, Phys. Letters **332**(1994), 349.
- [26] K. Morita, Prog. Theor. Phys. **96**(1996), 787.
- [27] K. Morita, Prog. Theor. Phys. **96**(1996), 801.
- [28] C. P. Martin, J. M. Gracia-Bondía and J. C. Várilly, The standard Model as a noncommutative geometry:the low energy region, hep-th/9605001.
- [29] B. Asquith, Phys. Letters **366**(1996), 220.
- [30] G. Landi, An Introduction to Noncommutative Spaces and Their Geometry, hep-th/9701078, Springer Verlag (1997).
- [31] L. Brink, J. H. Schwarz and J. Sherk, Nucl.Phys.**B121**(1977),77;
- [32] F. Gliozzi, J. Sherk and D. Olive, Nucl.Phys.**B122**(1977),253.
- [33] E. Alvarez, J. M. Gracia-Bondía and C. P. Martin, Phys.Lett.**306**(1993), 55;
- [34] E. Alvarez, J. M. Gracia-Bondía and C. P. Martin, Phys.Lett.**329**(1994), 259;
- [35] T. Shinohara, K. Nishida, H. Tanaka and I. Sogami, Prog.Theor.Phys.**96**(1996), 1179;
- [36] Y. Okumura, Prog.Theor.Phys.**98**(1997), 1333;hep-th/9707350, to appear in European Physical Journal **C**.
- [37] A. H. Chamseddine, A. Connes, hep-th/9606001, Comm. Math. Phys. **182**(1996), 155.
- [38] L. Calminati, B. Iochum, T. Schöcker, Noncommutative Yang-Mills and Noncommutative Relativity: A Bridge over Trouble Water, hep-th/9706105.